

A clinicopathological study of lesions of bone

Dnyanada Kokode*

Department of Pathology, NKP Salve Institute of Medical Sciences and Research Centre, Nagpur-440019, India

Abstract

Introduction: Bone tumours are neoplasms originating in the skeletal system that are within or closely related to the bone tissue. A spectrum of pathological bone lesions can present in any form from inflammatory to neoplastic conditions. They account for 0.2% of all tumours in humans.

Aims and objectives: To study the clinical and pathological spectrum of lesions of bone.

Material and methods: This study was carried out in the Department of Pathology, in a Tertiary health care hospital. It was a hospital based cross sectional study. After obtaining detailed clinical history and examination, biopsies and resected specimen were received in 10% formalin, gross findings were noted and histopathological examination was done.

Results: Histopathological evaluation was done in all 106 cases, obtained in a period of 2 years in the tertiary care hospital. In this study, non-neoplastic and benign neoplastic were the commonest bone lesions which accounted for 40.6% each, followed by malignant lesions 15.1% and metastatic lesions 3.7%. The maximum number of bone lesions occurred in second decade of life with a male to female ratio of 1.35:1. The most common presenting feature of all bone lesions was pain. The commonest site of all bone lesions in this study was lower end of femur followed by proximal end of tibia. The most common benign neoplastic neoplasms in this study was giant cell tumour followed by Osteochondroma. Among malignant neoplasm the most common was Osteosarcoma. Epithelial malignancies were the most common to metastasise to bone.

Conclusion: Histopathology is the gold standard for the precise diagnosis of the vast number of bone lesions. Since the exact diagnosis of bone tumours is at times difficult, a joint approach integrating clinical, radiological and histopathological findings is necessary to increase accuracy.

Introduction

Bone tumours are neoplasms originating in the skeletal system that are within or closely related to the bone tissue [1,2]. A spectrum of pathological bone lesions can present in any form from inflammatory to neoplastic conditions [1,2]. These lesions are diverse in their clinical and morphological features and range in behaviour from innocuous to rapidly fatal [3]. Accurate diagnosis, proper staging and appropriate treatment are thus necessary to ensure maximum patient survival and maintain optimal function of the affected body parts [4].

Primary bone tumours are relatively uncommon lesions [5]. They account for 0.2% of all tumours in humans and some of these tumours display marked inter- and intranational variations in incidence, site and age distribution [6]. Some relevant demographic features like age, gender and skeletal site are important factors while making a diagnosis [1,7,8]. The lesions predominantly occur in two age groups: adults over 40 years of age and children in the first and second decade of life. Various etiological agents including chemotherapy, radiation, trauma, infections and pre-existing bone lesions have been implicated. Common presentations are progressive pain, swelling, tenderness, restriction in joint mobility and pathological fracture in some of the cases [2,9].

In an Indian study, the most common bone involved was femur [4]. In chronic osteomyelitis and tuberculous lesions; femur and vertebrae respectively are known to be involved commonly. However, some benign processes such as osteomyelitis can mimic malignant tumours, and some malignant lesions such as metastases or myeloma, can mimic benign [3,4].

Aims and objectives

To study the clinical and pathological spectrum of lesions of bone.

Material and methods

This study was carried out in the Department of Pathology, in a Tertiary health care hospital after obtaining Institutional Ethics Committee Clearance. It was a hospital based cross sectional study. Patients presenting with lesions in bone of all age groups and both sexes were included in the research study. Detailed clinical history, radiological findings and other investigations were noted. FNAC was done wherever possible. All bone biopsies and resected specimens received during a period of 2 years were studied. Biopsies and resected specimen were received in 10% formalin and gross findings were noted. Tissues were processed and stained with routine haematoxylin and eosin stain. Histopathological examination was done under light microscope.

Results

Histopathological evaluation was done in all 106 cases, obtained in a period of 2 years in the tertiary care hospital. In this study, non-neoplastic and benign neoplastic were the commonest bone lesions which accounted for 40.6 % each, followed by malignant lesions 15.1% and metastatic lesions 3.7%.

***Correspondence to:** Dnyanada Kokode, Department of Pathology, NKP Salve Institute of Medical Sciences and Research Centre, Nagpur-440019, India, E-mail: dr.radhakokode@gmail.com

Received: February 06, 2019; **Accepted:** February 20, 2019; **Published:** February 25, 2019

In this study the age of patients varied from 9 years to 81 years. The mean age was 29.72 years. There was an overall male preponderance with a male to female ratio of 1.35:1. The maximum number of bone lesions occurred in the age group of 11-20 years followed by 21-30 years of age group as mentioned in (Table 1).

The most common presenting feature of all bone lesions was pain (96 cases) followed by joint movement restriction (89 cases). The commonest site of all bone lesions in this study was lower end of femur with 25 cases (30.49%) followed by proximal end of tibia (20 .73%) (Table 2). Among Non-neoplastic lesions, Chronic Osteomyelitis (32 cases, 33.9%) were most common followed by Tuberculosis Osteomyelitis (5 cases, 4.7%) (Figure 1a and 1b).

The most common benign neoplasm was Giant cell tumour (10 cases). The most common malignant neoplasm was Osteosarcoma (7 cases).

Discussion

Histopathological examination is the “gold standard” for definite diagnosis of bone lesions [1]. This study describes the pattern and

frequencies of bone lesions including non- neoplastic (infectious), benign and malignant neoplasms.

The benign lesions (including non-neoplastic and neoplastic) were 86 cases (81.2%) and malignant (primary and secondary) were 20 cases (18.8%) This data correlated with the other studies mentioned in (Table 3) [1,3,4,10,11].

In this study the age of patients varied from 9 years to 81 years. The mean age was 29.72 years. There was an overall male preponderance with a male to female ratio of 1.35:1. These results were comparable to various studies as mentioned in (Table 4). Similar results were reported in studies done by [1,3,4,11] Jain S, *et al.* found a male to female ratio of 0.53:1.

The second decade was the commonest age group involved in our study, which correlated with other studies [1,3,4,11-15].

The commonest site of all bone lesions in this study was lower end of femur, 25 cases (30.49%) followed by proximal end of tibia (20 .73%), which parallels with the results of other studies [1,4,11-15].

Table 1. Gender wise distribution of bone Lesions

Sr. No.	Histological Type	Males	Females	Total	M:F
I.	Infectious etiology				
1.	Subacute pyogenic osteomyelitis	1	1	2	1:1
2.	Chronic osteomyelitis	25	11	32	2.3:1
3.	Tuberculous osteomyelitis	3	2	5	1.5:1
II.	CHONDROGENIC TUMORS				
1.	Benign				
	a. Osteochondroma	5	2	7	2.5:1
	b. Chondroma	2	2	4	1:1
2.	Intermediate				
	a. Chondroblastoma	0	3	3	0:3
3.	Malignant				
	a. Chondrosarcoma	1	2	3	1:2
III.	OSTEOGENIC TUMORS				
1.	Benign				
	a. Osteoid osteoma	2	2	4	1:1
2.	Intermediate				
	a. Osteoblastoma	0	2	2	0:2
3.	Malignant				
	a. Osteosarcoma	4	3	7	1.3:1
IV.	FIBROGENIC TUMOURS				
	Fibroma of bone	1	1	2	1:1
V	EWINGS SARCOMA	2	1	3	2:1
VI.	HAEMATOPOIETIC NEOPLASMS				
1.	Plasma cell myeloma	2	0	2	2:0
VII.	GIANT CELL TUMOURS				
	Giant cell tumour	5	5	10	1:1
VIII	NOTOCHORDAL TUMOURS				
I	Malignant				
	Chordoma	2	1	3	2:1
IX	MISCELLANEOUS LESIONS				
1.	Benign				
	Osteofibrous dysplasia	1	2	3	1:2
2.	Intermediate				
	Aneurysmal bone cyst	2	1	3	2:1
	Langerhans cell histiocytosis	0	1	1	0:1
X	MISCELLANEOUS TUMOURS				
	Adamantinoma	0	2	2	0:2
XI.	Metastasis	1	3	4	1:3
	Total	61 (57.5%)	45 (42.5%)	106 (100%)	1.35:1

Table 2. Common presentation of bone lesions




Sr No.	Histological types	Pain (%)	Swelling (%)	Sinus/ Ulcers (%)	Fracture (%)	Trauma (%)	Joint movement Restriction (%)
1.	Osteomyelitis	32(100)	23(72)	2(6.3)	7(22)	11(34)	28(87.5)
2.	Tuberculous osteomyelitis	5 (100)	4(80)	1(20)	2(40)	0	4(80)
3.	Benign chondrogenic tumors	11(100)	10(91)	0	0	1(0.9)	10(91)
4.	Intermediate chondrogenic tumors	3(100)	3(100)	0	1(33.3)	1(33.3)	3(100)
5.	Malignant chondrogenic tumors	1(33.3)	3(100)	0	0	0	3(100)
6.	Benign osteogenic tumors	5(100)	4(80)	0	0	0	4(80)
7.	Intermediate osteogenic tumors	2(100)	2(100)	0	0	0	2(100)
8.	Malignant osteogenic tumors	7(100)	6(86)	2(28.6)	1(14.3)	1(14.3)	6(86)
9.	Fibroma of bone	2(100)	1(50)	0	0	0	2(100)
10.	Plasma cell myeloma	2(100)	1(50)	0	0	0	2(50)
11.	Giant cell tumour	8(80)	9(90)	0	0	2(20)	9(90)
12.	Chordoma	3(100)	2(66.6)	0	0	0	2(66.6)
13.	Osteofibrous dysplasia	1(33.3)	3(100)	0		1(33.3)	3(100)
14.	Aneurysmal bone cyst	2(66.6)	3(100)	0	0	0	2(66.6)
15.	Langerhans cell histiocytosis	1(100)	1(100)	0	0	1(100)	1(100)
16.	Ewings sarcoma	3(100)	3(100)	0	0	0	2(66.6)
17.	Adamantinoma	2(100)	2(100)	0	0	0	2(100)
18.	Metastasis	4(100)	1(25)	0	3(75)	1(25)	4(100)
	Total	96	82	5	14	19	89

Table 3. Distribution of bone tumours according to WHO classification

Sr. No.	Histological Type	Number	Frequency (%) in the total bone lesions in our study
I.	CHONDROGENIC TUMORS		
1.	Benign		
	a. Osteochondroma	7	6.6 %
	b. Chondroma	4	3.8 %
2.	Intermediate		
	a. Chondroblastoma	3	2.8 %
3.	Malignant		
	a. Chondrosarcoma	3	2.8 %
II.	OSTEOGENIC TUMORS		
1.	Benign		
	a. Osteoid osteoma	4	3.8 %
2.	Intermediate		
	a. Osteoblastoma	2	1.9 %
3.	Malignant		
	a. Osteosarcoma	7	6.6 %
III	FIBROGENIC TUMORS		
	Fibroma of bone	2	1.9 %
IV.	EWINGS SARCOMA	3	2.8 %
V.	HAEMATOPOIETIC NEOPLASMS		
1.	Plasma cell myeloma	2	1.9 %
VI.	GIANT CELL TUMOURS		
	Giant cell tumour	10	9.4 %
VII.	NOTOCHORDAL TUMOURS		
1	Malignant		
	a. Chordoma	3	2.8 %
VIII	MISCELLANEOUS TUMOURS		
1.	Benign		
	a. Osteofibrous dysplasia	3	2.8 %
2.	Intermediate		
	a. Aneurysmal bone cyst	3	2.8 %
	b. Langerhans cell histiocytosis	1	0.9 %
IX	MISCELLANEOUS LESIONS		
	a. Adamantinoma	2	1.9 %
X	Metastasis	4	3.7 %
	Total	63	59.4 %

Table 4. Frequency of benign and malignant tumours in various studies

Sr. No	Studies	Giant cell tumor commonest benign neoplastic lesion	Osteochondroma 2 nd commonest benign neoplastic lesion	Number of cases of Osteosarcoma (commonest bone malignancy)
1	S Bamanikar et al, 2015 [1]	16 (19.51%)	5 (21.42%)	7 (8.53%)
2	S Jain et al, 2011 [15]	24 (20.51%)	26(22.22%)	13(35.14%)
3	Abdulkarem FB et al, 2009 [11]	18 (23.4%)	6(33.3%)	7(9.1%)
4	B Wamisho et al, 2010 [4]	22 (10.73%)	22(10.73%)	45(21.95%)
5	Present Study	10 (9.43 %)	7 (6.9 %)	7 (6.9%)

**Figure 1.** Tuberculous Osteomyelitis**Figure 1a.** X ray spine showing blurred paradiscal margins and sclerosis of vertebral bodies**Figure 1b.** Photomicrograph showing Tuberculous lesion of bone showing well formed granulomas. (Hematoxylin and eosin stain 10X)**Figure 2.** Osteosarcoma**Figure 2a.** X ray showing mixed lytic and blastic lesion at upper end of tibia**Figure 2b.** Gross image showing growth involving upper end of tibia**Figure 2c.** Photomicrograph showing abundant osteoid production along with hyaline cartilage. (Hematoxylin and eosin stain 40x)**Figure 3.** Metastatic deposits of malignancy**Figure 3a.** MRI spine showing sclerosis and lytic lesion at T12 vertebrae**Figure 3b.** Photomicrograph showing metastasis of invasive ductal carcinoma of breast in bone. (Hematoxylin and eosin stain 40x)

The most common benign tumour in this study was giant cell tumour (10 cases, 9.43%) followed by Osteochondroma (7 cases, 6.6%). We had 10 (9.43%) cases of giant cell tumours in our study. Most of the cases were seen in 11-20 years of age with Male to female ratio of 1:1. The most common site of presentation was radius (5 cases, 4.7%). Most of the lesions presented radiologically as a lytic lesion in the bone with surrounding reactive bone. These findings correlated with the findings of above reference studies.

In our study, malignant lesions included 7 cases (6.9%) of Osteosarcoma, 3 cases (2.8%) of Chondrosarcoma and Ewing's sarcoma each. Most of the cases of Osteosarcoma were reported in 11- 20 years of age. The commonest site was femur. In other studies, it ranged in frequency with osteosarcoma ranging from 8.53% to 35.14% (Figure 2a,2b and 2c). Patients with Osteosarcoma generally present to the hospital very late (average 4–5 months) after onset of symptoms. Increased awareness and improved referral systems may decrease this delay and can help for proper treatment [4].

Of the 4 cases of metastatic lesions in our study, epithelial malignancies were most common (3 cases, 2.9%) to metastasize followed by soft tissue malignancy (Figure 3a,b). The primary sites for the epithelial metastasis were from breast, colon and kidney. The prevalence of multiple primary epithelial malignant neoplasms varies from 0.734% to 11.7%. The case of metastatic soft tissue tumour was of, 13 year old female patient who presented with difficulty in walking. Patient had pathological fracture neck femur and greater trochanter. Biopsy revealed features of Liposarcoma. Extensive immunohistochemical staining was performed. The tumour cells were immune-negative for S-100 protein, CDK- 4, CD 68, CD 163, Cytokeratin, EMA, Melan A, LCA, Desmin, SMA, Calretinin, Inhibin, Synaptophysin and Chromogranin A. Thus ruling out other differential diagnosis, the features were compatible with a pleomorphic liposarcoma metastasizing to the bone.

Conclusion

In conclusion, this study showed that primary bone lesions were mainly benign, occurred predominantly in the second decade of life with a male preponderance. Chronic Osteomyelitis, Giant cell tumour and Osteosarcoma were the most common non-neoplastic, benign and primary malignant bone lesions, respectively. Epithelial malignancies were the most common primary tumours to metastasize to bone. Since the exact diagnosis of bone tumours is at times difficult,

a joint approach integrating clinical, radiological and histopathological findings is necessary to increase accuracy [4,10]. Histopathology is the gold standard for the precise diagnosis of the vast number of bone lesions [3,12,13]. If the diagnosis is not confirmed by histology, then there is a risk of inappropriate surgery being carried out [11,14]. The clinical and pathologic spectrum of bone lesions seen at our centre were similar to those reported from other national and international studies.

References

1. Bamanikar S, Pagaro P, Kaur P, Chandanwale S, Bamanikar A, et al. (2015) Histopathological Study of Primary Bone Tumours and Tumour-Like Lesions in a Medical Teaching Hospital. *JKIMSU* 4: 46-55.
2. Björn R (1994) Bone Embryology. *Anat Embryol* 189: 19-24.
3. Deoghare SB, Prabhu MH, Ali SS, Inamdar SS (2017) Histomorphological Spectrum of Bone Lesions at Tertiary Care Centre *Int. J. Life. Sci. Scienti. Res* 3: 980-985.
4. Wamisho B, Admasie D, Negash B, Tinsay M (2009) Osteosarcoma of limb bones: a clinical, radiological and histopathological diagnostic agreement at Black Lion Teaching Hospital, Ethiopia. *Malawi Med J* 21: 62-65. [[Crossref](#)]
5. Johnson LC (1953) A general theory of bone tumors. *Bull NY Acad Med* 29: 164-171. [[Crossref](#)]
6. Moodie RL (1923) Paleopathology. An Introduction to the Study of Ancient Evidences of Disease. pp: 567-574.
7. Saanna, Bovee J, Hornick J, Lazar A (2013) WHO classification of tumors of Soft Tissue and bone 4: 5-7.
8. Meltzer PS, Kallioniemi A, Trent JM (2002) History of cancers 93: 113-115.
9. Peltier LF (1985) Historical note on bone and soft tissue sarcoma. *J Surg Oncol* 30: 201-205. [[Crossref](#)]
10. Chaurasia BD (2013) Human Anatomy: Regional and applied dissection and clinical. 7: 2-9.
11. Abdulkareem FB, Eyesan SU, Akinde OR, Ezembakwe ME, Nnodu OE (2007) Pathological study of Bone Tumours at the National Orthopaedic Hospital, Lagos, Nigeria. *West African J Medicine* 26: 306-11.
12. Singh M, Singh S, Jain J, Singh KT (2010) Chronic suppurative osteomyelitis of maxilla mimicking actinomycotic osteomyelitis: A rare case report. *Natl J Maxillofac Surg* 1:153-156. [[Crossref](#)]
13. Marx R (1991) Chronic osteomyelitis of the jaws. *Oral Maxillofac Clin North Am* 14: 3: 367-381.
15. Jones A, Isaacs D (2013) Systematic review of duration and choice of systemic antibiotics therapy for acute haematogenous bacterial osteomyelitis in children. *J Paediatr Child Health* 49: 760-768.
16. Jain KS, Ravishankar RM, Rupakumar CS, Gadiyar HB, Manjunath GV (2011) Bone tumors in a tertiary care hospital of south India: A review 117 cases. *IJMPO* 32: 82-85.