

## Fetal sacrococcygeal teratoma

Belliemi CV<sup>\*1</sup>, Molinaro F<sup>2</sup>, Sica M<sup>2</sup>, Angotti R<sup>2</sup>, Messina M<sup>2</sup>, Bocchi C<sup>3</sup>, Severi F<sup>3</sup> and Carbone S<sup>4</sup>

<sup>1</sup>Department of Mother and Infant, University Hospital, Siena, Italy

<sup>2</sup>Division of Pediatric Surgery, Department of Medical, Surgical and Neurological Sciences, University of Siena, Italy

<sup>3</sup>Division of Obstetrics, University of Siena, Italy

<sup>4</sup>Institute of radiology, University Hospital, Siena, Italy

An anterior myelomeningocele had been suspected in a female fetus on the basis of fetal ultrasonography; prenatal MRI at the 24<sup>th</sup> week of gestation confirmed a cystic mass in the anterior sacrococcygeal

area [1]. The mass was surgically removed at one month of age, and hystopathological analysis determined it was a mature teratoma (Figure 1).



Figure 1. Prenatal diagnosis (fetal MRI): a tumor in the fetal sacrococcygeal area.

**Copyright:** ©2017 Bellieni CV. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Correspondence to:** Carlo Valerio Bellieni, Department of Mother and Infant, University Hospital, Siena, Italy, E-mail: cvbellieni@gmail.com

**Received:** May 04, 2017; **Accepted:** May 17, 2017; **Published:** May 20, 2017