

Attenuation of high sucrose diet–induced insulin resistance in ABC transporter deficient *white* mutant of *Drosophila melanogaster*

Valeriya Navrotskaya¹, Gregory Oxenkrug^{2*}, Lyudmila Vorobyova¹ and Paul Summergrad²

¹Department of Genetics and Cytology, V.N. Karazin Kharkiv National University, Ukraine

²Psychiatry and Inflammation Program, Department of Psychiatry, Tufts University/Tufts Medical Center, USA

Abstract

Exposure to high sugar diet (HSD) is an experimental model of insulin resistance (IR) and type 2 diabetes (T2D) in mammals and insects. In *Drosophila*, HSD-induced IR delays emergence of pupae from larvae and eclosion of imago from pupae. Understanding of mechanisms of IR/T2D is essential for refining T2D prevention and treatment strategies. Dysregulation of tryptophan (Trp)-kynurenine (Kyn) pathway was suggested as one of the mechanisms of IR/T2D development. Rate-limiting enzyme of Trp-Kyn pathway in *Drosophila* is Trp 2,3-dioxygenase (TDO), an evolutionary conserved ortholog of human TDO. We previously reported attenuation of HSD-induced IR in *vermilion* mutants with inactive TDO. Conversion of Trp to Kyn is regulated not only by TDO activity but by intracellular Trp transport via ATP-binding cassette (ABC) transporter encoded by *white* gene in *Drosophila*. In order to evaluate the possible impact of deficient intracellular Trp transport on the inducement of IR by HSD, we compared the effect of HSD on pre-imago development in wild type flies, Canton-Special (C-S), and C-S flies containing *white* gene, *white* (C-S). Presence of *white* gene attenuated (by 50%) HSD-induced delay of pupae emergence from larvae and female and male imago eclosion from pupae. Present study together with our earlier report reveals that both decreased TDO activity (due to *vermilion* gene mutation) or deficient Trp transport into cell without affecting TDO levels (due to *white* gene mutation) attenuate HSD-induced development of IR in *Drosophila* model of T2D. Our data provide further support for hypothesis that dysregulation of Trp-Kyn pathway is one of the pathophysiological mechanisms and potential target for early diagnosis, prevention and treatment of IR/T2D.

Introduction

Diabetes mellitus is a public health problem, which affects a millions worldwide. Most diabetes cases are classified as type 2 diabetes mellitus (T2D), which is highly associated with obesity [1]. Impaired glucose tolerance (pre-diabetes) is a high-risk factor for T2D [2]. Insulin resistance (IR) is associated with prediabetes before T2D could be diagnosed. Further elucidation of IR mechanisms is needed for prevention and treatment of T2D. It was suggested that dysregulation of kynurenine (Kyn) pathway of tryptophan (Trp) metabolism is one of the mechanisms of development of IR/T2D [3-8]. We found IR correlation with activity of Trp-Kyn pathway in Hepatitis C virus (HCV) patients [9], and elevation of plasma concentrations of Kyn downstream metabolites in T2D patients [10]. Experimental model of T2D was established in *Drosophila melanogaster* [11]. There are four distinct stages in the life of *flies*: egg, larva, pupa, and imago (adult). High sugar diet (HSD) induces IR in larva and T2D in imago [12]. IR delays emergence of pupae from larva and imago eclosion from pupae [13]. In *Drosophila*, tryptophan 2,3-dioxygenase (TDO), enzyme, catalyzing formation of Kyn (via N-formyl-kynurenine) from Trp, is encoded by *vermilion* gene [14]. We found attenuation of HSD-induced development of IR in *vermilion* mutants with inactive TDO [15]. TDO is substrate-activated intracellular enzyme. Therefore, conversion of Trp to Kyn is regulated not only by TDO activity but by Trp transport into cells as well [16]. Import of Trp into cell is mediated by ATP-binding cassette (ABC) transporters encoded by *white* gene in *Drosophila*. *White* mutations do not alter levels of TDO, but interfere with the ability of cells to take up Trp [17]. Therefore, mutations of both

TDO (*vermilion*) and ABC transporter (*white*) genes have a similar effect, *i.e.*, decreased formation of Kyn from Trp. In order to evaluate the possible impact of impairment of intracellular Trp transport on the inducement of IR by HSD, we compared the effect of HSD on pre-imago development in wild type flies, Canton-Special (C-S), and C-S flies containing *white* gene, *white* (C-S).

Materials and methods

Flies from C-S the collection of V.N. Karazin Kharkiv National University were used in the study. Wild type *Drosophila melanogaster* and *white* (C-S) mutants flies were maintained at 23°C in a 12:12 light: dark period on a standard nutrition medium consisting of sugar, yeast, agar and semolina. Experimental eggs were obtained from parents with synchronized egg laying. Before eggs laying sucrose (0.67M) was added to nutrition medium. Emerging time was taken as the period from the time of synchronized egg laying to the time of larvae emergence into pupae as described elsewhere [15]. Appearance of female and male

Correspondence to: Gregory F. Oxenkrug, Tufts Medical Center, 800 Washington St, #1007, Boston, MA, 02111, USA, **E-mail:** goxenkrug@tuftsmedicalcenter.org

Key words: *white* mutants, ABC-transporter, *drosophila*, high sugar diet, insulin resistance, type 2 diabetes, kynurenine, tryptophan, obesity

Received: January 10, 2016; **Accepted:** February 04, 2016; **Published:** February 08, 2016

imago from pupae was recorded as time of eclosion. The study was carried out in November and December 2015.

Statistical analysis

Data were expressed as mean±standard deviation (hours of pupae emergence and imago eclosion). Differences between experimental groups were evaluated by Mann Whitney, two-tailed test.

Results

Pupae emergence from larva

Emergence time of C-S wild type flies maintained on standard nutrition medium was 7% shorter than emergence time of *white* (C-S) mutants ($p<0.0001$) (Table 1). HSD delayed pupae emergence from larva of wild type flies, C-S, in comparison with flies maintained on standard nutrition medium by almost 3 days (40%). In *white* (C-S) mutants HSD delayed pupae emergence from larvae by 1.6 days (20%) in comparison with flies maintained on standard nutrition medium. Therefore, presence of *white* gene attenuated HSD-induced delay of pupae emergence from larvae by 50% (Table 1).

Imago eclosion from pupae

There was no significant gender effect on imago eclosion time (Table 2). Eclosion times of both female and male C-S wild type flies maintained on standard nutrition medium were about 8% shorter than eclosion times of *white* (C-S) mutants ($p<0.0001$). HSD delayed imago eclosion in C-S flies in comparison with flies maintained on standard nutrition medium by 2.75 days (20.5%) in females and by 2.60 days (19.8%) in males. HSD delayed imago eclosion in *white* (C-S) mutants by 1.3 days (9%) days in females and by 1 day (7%) male flies. Therefore, presence of *white* gene attenuated HSD-induced imago eclosion by 50% in comparison with wild type flies (Table 2).

Discussion

Present data indicate that HSD delays pupae emergence from larva and imago eclosion in wild type C-S flies in accordance with literature data [11-13]. We are not aware of studies of the effect of HSD on preimaginal stages (from egg through larva and pupa to imago) of ABC transporter deficient *white* flies. Notably, the emergence time of *white* (C-S) mutants maintained on standard nutrition medium

was longer than emergence time of C-S wild type flies. Similar results were seen for imago eclosion time in *white* (C-S) and Canton-S flies. This observation may suggest that development rate of egg to adult fly is naturally delayed in mutant flies as compared to wild flies. We observed that HSD reversed this tendency in mutants in contrast to HSD-induced delay of preimaginal development in wild flies: HSD-induced delay of pre-imaginal development was two times shorter in *white* (C-S) mutants than in wild type C-S flies. Considering that HSD-induced delay of preimaginal development is caused by IR [11-13], our data suggest that mutation of *white* gene attenuates HSD-induced IR.

Attenuation of HSD-induced IR in *white* (C-S) flies most likely depends on deficiency of ABC transporter that mediates Trp import into cell, and, thus, decreases availability of Trp, a substrate for intracellularly located TDO [16]. Decreased availability of Trp as a substrate for TDO results in decreased Kyn formation from Trp, without affecting TDO activity [17]. TDO-regulated KYN formation from Trp begins at the end of the third larval instar in the cells of the anterior region of fat body (analog of liver and fat tissues in humans) [14]. Immediate metabolic response to HSD in *Drosophila* is an increase of glucose in hemolymph [12]. Hyperglycemia induces production of O_2^- and H_2O_2 , and generation of mitochondrial reactive oxygen species in rodents [18], and activates hypothalamic-pituitary-adrenal axis and, consequently, increases secretion of cortisol, a hormonal inducer of TDO, in rat model of T2D [19] and in T2D patients [20]. Notably, high glucose selectively inhibits ABC transporter G1 subtype, involved in Trp transport, in human macrophages [21]. Therefore, mutation of *white* gene, that prevents Trp import into cell, might confer resistance to HSD-induced IR and T2D, an aging associated disease [4]. Notably, we reported pronged life span in *white* mutants [22] while resistance to high glucose-induced oxidative stress was associated with longevity in rodents [18]. Considering that *white* mutants have deficient transport of both Trp and guanine, attenuation of HSD-induced IR might depend not only on Trp but guanine transporter deficiency as well. While such a possibility could not be ruled out based on available data, some other factors might affect functions of *white* mutant, the parent strain of *Methuselah* *Drosophila* [23,24]. Further studies, e.g., of the effect of ABC-transporter inhibitor, 5-methylTrp, [25] on HSD-induced IR might help to differentiate between impairment of guanine- and Trp-ABC transporters.

Table 1. Effect of high sucrose diet on time of pupae emergence from larvae in *white* (C-S) and Canton-S flies.

Genotype and treatment	Standard Diet	High Sucrose Diet	Delay of emergence (%)
Canton-S	182.9 ± 18.9 (n=516)	253.8 ± 24.5* (n=465)	40
<i>white</i> (C-S)	195.4 ± 22.2* (n=385)	234.2 ± 31.6** (n=261)	20

Mean ± standard deviation (hrs); n: number of pupae

Mann-Whitney two tailed test: * $p=0.0001$ in comparison with Canton-S; ** $p=0.0001$ in comparison with Standard diet.

Table 2. Effect of high sucrose diet on imago eclosion *white* (C-S) and Canton-S flies.

Genotype and treatment	Standard Diet	High Sucrose Diet	Delay of eclosion (%)
Canton-S:			
Female	322.0 ± 17.1 (n=144)	388.1 ± 23.9* (n=144)	20.5
Male	327.8 ± 16.7 (n=126)	392.7 ± 24.1* (n=111)	19.8
<i>white</i> (C-S):			
Female	349.3 ± 36.3 ** (n=165)	380.5 ± 32.1* (n=108)	9
Male	357.9 ± 35.2 ** (n=163)	382.7 ± 33.6* (n=114)	7

Mean ± standard deviation (hrs); n: number of imago.

Mann-Whitney two tailed test: * $p=0.0001$ in comparison with flies on standard diet; ** $p=0.0001$ in comparison with Canton-S

Considering high association of obesity and T2D [26] and possible involvement of Trp-Kyn pathway in development of obesity [3,6,27-29], it might be of interest to study development of HSD-induced obesity [11 *white* mutants of *Drosophila melanogaster*.

In conclusion, present study revealed that attenuation of HSD-induced development of IR in *Drosophila* model of T2D might result from the deficiency of Trp transport into cell without affecting TDO levels. We have previously reported that TDO deficiency attenuated HSD-induced IR development as well [15]. Our data provide further support for hypothesis that dysregulation of Trp-Kyn pathway is one of the pathophysiological mechanisms and potential target for early diagnosis, prevention and treatment of IR/T2D [4-7].

Acknowledgements

GF Oxenkrug is a recipient of MH104810. Paul Summergrad is a non-promotional speaker for CME outfitters, Inc.

References

1. Tao Z, Shi A, Zhao J (2015) Epidemiological Perspectives of Diabetes. *Cell Biochem Biophys* 73: 181-185. [Crossref]
2. Herder C, Nuotio ML, Shah S, Blankenberg S, Brunner EJ, et al. (2014) Genetic determinants of circulating interleukin-1 receptor antagonist levels and their association with glycemic traits. *Diabetes* 63: 4343-4359. [Crossref]
3. Connick JH, Stone TW (1985) The role of kynurenines in diabetes mellitus. *Med Hypotheses* 18: 371-376. [Crossref]
4. Oxenkrug GF (2010) Metabolic syndrome, age-associated neuroendocrine disorders, and dysregulation of tryptophan-kynurenine metabolism. *Ann NY Acad Sci* 1199: 1-14. [Crossref]
5. Oxenkrug G (2013) Insulin Resistance and Dysregulation of Tryptophan-Kynurenine and Kynurenine Nicotinamide Adenine Dinucleotide Metabolic Pathways. *Mol Neurobiol* 48: 294-301. [Crossref]
6. Oxenkrug G (2015) 3-hydroxykynurenine acid and type 2 diabetes: implications for aging, obesity, depression, Parkinson's disease and schizophrenia. In: *Tryptophan Metabolism: Implications for Biological Processes, Health and Diseases, Molecular and Integrative Toxicology* (A. Engin, A.B. Engin - eds), pp.173-195.
7. Oxenkrug GF (2015) Role of kynurenine pathway in insulin resistance: towards kynurenine hypothesis of insulin resistance and diabetes. Springer International Publishing, Switzerland pp: 169-178. [Crossref]
8. Pedersen ER, Tuseth N, Eussen SJ, Ueland PM, Strand E, et al. (2015) Associations of plasma kynurenines with risk of acute myocardial infarction in patients with stable angina pectoris. *Arterioscler Thromb Vasc Biol* 35: 455-462. [Crossref]
9. Oxenkrug GF, WA Turski, W Zgrajka, JV Weinstock, P Summergrad (2013) Tryptophan-Kynurenine Metabolism and Insulin Resistance in Hepatitis C Patients. *Hepatitis Research and Treatment 2013*: 149247.
10. Oxenkrug GF (2015) Increased Plasma Levels of Xanthurenic and Kynurenic Acids in Type 2 Diabetes. *Mol Neurobiol* 52: 805-810. [Crossref]
11. Musselman LP, Fink JL, Narzinski K, Ramachandran PV, Hathiramani SS, et al. (2011) A high-sugar diet produces obesity and insulin resistance in wild-type *Drosophila*. *Dis Model Mech* 4: 842-849. [Crossref]
12. Pasco MY, Léopold P (2012) High sugar-induced insulin resistance in *Drosophila* relies on the lipocalin Neural Lazarillo. *PLoS One* 7: e36583. [Crossref]
13. Rovenko BM, Perkhulyn NV, Gospodaryov DV, Sanz A, Lushchak OV, et al. (2015) High consumption of fructose rather than glucose promotes a diet-induced obese phenotype in *Drosophila melanogaster*. *Comp Biochem Physiol A Mol Integr Physiol* 180: 75-85. [Crossref]
14. Rizki TM, Rizki RM (1964) Factors affecting the intracellular synthesis of kynurenine. *J Cell Biol* 21: 27-33. [Crossref]
15. Navrotskaya V, Oxenkrug G, Vorobyova L, P Summergrad (2015) Attenuation of high sucrose diet-induced insulin resistance in tryptophan 2,3-dioxygenase deficient *Drosophila melanogaster*. *Integr Obes Diabetes* 1: 93-95. [Crossref]
16. Oxenkrug GF (2007) Genetic and hormonal regulation of tryptophan kynurenine metabolism: implications for vascular cognitive impairment, major depressive disorder, and aging. *Ann NY Acad Sci* 1122: 35-49. [Crossref]
17. Ewart GD, Howells AJ (1998) ABC transporters involved in transport of eye pigment precursors in *Drosophila melanogaster*. *Methods Enzymol* 292: 213-224. [Crossref]
18. Labinskyy N, Mukhopadhyay P, Toth J, Szalai G, Veres M, et al. (2009) Longevity is associated with increased vascular resistance to high glucose-induced oxidative stress and inflammatory gene expression in *Peromyscus leucopus*. *Am J Physiol Heart Circ Physiol* 296: H946-956. [Crossref]
19. Elahi-Moghaddam Z, Behnam-Rassouli M, Mahdavi-Shahri N, Hajinejad-Boshroue R, Khajouee E (2013) Comparative study on the effects of type 1 and type 2 diabetes on structural changes and hormonal output of the adrenal cortex in male Wistar rats. *J Diabetes Metab Disord* 12: 9. [Crossref]
20. Chiodini I, Torlontano M, Scillitani A, Arosio M, Bacci S, et al. (2005) Association of subclinical hypercortisolism with type 2 diabetes mellitus: a case-control study in hospitalized patients. *Eur J Endocrinol* 153: 837-844. [Crossref]
21. Mauerer R, Ebert S, Langmann T (2009) High glucose, unsaturated and saturated fatty acids differentially regulate expression of ATP-binding cassette transporters ABCA1 and ABCG1 in human macrophages. *Exp Mol Med* 41: 126-132. [Crossref]
22. Oxenkrug GF (2010) The extended life span of *Drosophila melanogaster* eye-color (white and vermilion) mutants with impaired formation of kynurenine. *J Neural Transm (Vienna)* 117: 23-26. [Crossref]
23. Borycz J, Borycz JA, Kubo'w A, Lloyd V, Meinertzhagen IA (2008) *Drosophila* ABC transporter mutants white, brown and scarlet have altered contents and distribution of biogenic amines in the brain. *J Exp Biol* 211: 3454-3466. [Crossref]
24. Petrosyan A, Hsieh IH, Saberi K (2007) Age-dependent stability of sensorimotor functions in the life-extended *Drosophila* mutant methuselah. *Behav Genet* 37: 585-594. [Crossref]
25. Oxenkrug GF, Navrotskaya V, Vorobyova L, Summergrad P (2011) Extension of life span of *Drosophila melanogaster* by the inhibitors of tryptophan-kynurenine metabolism. *Fly (Austin)* 5: 307-309. [Crossref]
26. Gallagher EJ, LeRoith D (2015) Obesity and Diabetes: The Increased Risk of Cancer and Cancer-Related Mortality. *Physiol Rev* 95: 727-748. [Crossref]
27. Mangge H, Summers KL, Meinitzer A, Zelzer S, Almer G, et al. (2014) Obesity-related dysregulation of the tryptophan-kynurenine metabolism: role of age and parameters of the metabolic syndrome. *Obesity (Silver Spring)* 22: 195-201. [Crossref]
28. Wolowczuk I, Hennart B, Leloire A, Bessede A, Soichot M, et al. (2012) Tryptophan metabolism activation by indoleamine 2,3-dioxygenase in adipose tissue of obese women: an attempt to maintain immune homeostasis and vascular tone. *Am J Physiol Regul Integr Comp Physiol* 303: R135-143. [Crossref]
29. Favennec M, Hennart B, Caiazza R, Leloire A, Yengo L, et al. (2015) The kynurenine pathway is activated in human obesity and shifted toward kynurenine monoxygenase activation. *Obesity (Silver Spring)* 23: 2066-2074. [Crossref]

Copyright: ©2015 Navrotskaya V. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.